

BstP I

(BstE II, EcoO65 I)

G | G T N A C C
C C A N T G | G

Code No. 1025A **Size:** **2,000 U**
Conc.: **10 U/μl**

Supplied Reagents:
10X H Buffer **1 ml**
10X Loading Buffer **1 ml**

Storage Buffer: 10 mM Tris-HCl, pH 7.5
100 mM KCl
0.1 mM EDTA
1 mM DTT
0.15% Triton X-100
0.01% BSA
50% Glycerol

Storage: -20°C

Source: *Bacillus stearothermophilus*

General Reaction Mixture:
BstP I 1 μl
10X H Buffer 2 μl
Substrate DNA ≤ 1 μg
Sterile purified water up to 20 μl

Reaction Temperature: 60°C

Unit definition:

One unit is defined as the amount of this enzyme required to digest completely 1 μg of λ DNA in 50 μl of the reaction mixture at 60°C for 1 hr.

Quality Control Data:

Please see the Certificate of Analysis (CoA) for each lot. You can download the CoA on Takara Bio website.

Relative Activity in Takara Bio's Universal Buffers:

Universal Buffer	L	M	H	K	T (+BSA)
Relative Activity (%)	(<20)	(60)	100	(100)	(100)

(): Weak star activity is detected.

Ionic Effect on Activity in Basal Buffer:

Salt (mM)	0	20	50	80	100	150	200
NaCl (%)	20	30	60	70	100	50	20
KCl (%)	20	20	60	70	100	50	10

Composition of Basal Buffer:

50 mM Tris-HCl, pH 8.0
7 mM MgCl₂
100 mM NaCl
7 mM 2-mercaptoethanol

Number of Cleavage Sites in DNA:

		SV	φX	pBR	pUC	pUC	M13	Col
λ	Ad2	40	174	322	19	119	mp18	E1
13	10	0	0	0	0	0	0	2

Effect of DNA Methylation:

Enzyme activity is not affected by CG methylase.

Star activity:

Unrelated site may often be cut in the presence of high concentration of glycerol, and at low ionic strength.

Compositions of Universal Buffer (Stored at -20°C):

1. 10X L	100 mM Tris-HCl, pH7.5	4. 10X K	200 mM Tris-HCl, pH8.5
	100 mM MgCl ₂		100 mM MgCl ₂
	10 mM Dithiothreitol		10 mM Dithiothreitol
2. 10X M	100 mM Tris-HCl, pH7.5		1,000 mM KCl
	100 mM MgCl ₂	5. 10X T	330 mM Tris-Ac, pH7.9
	10 mM Dithiothreitol	(BSA-free)	100 mM Mg-Ac
	500 mM NaCl		5 mM Dithiothreitol
3. 10X H	500 mM Tris-HCl, pH7.5		660 mM K-Ac
	100 mM MgCl ₂		6. 0.1% BSA
	10 mM Dithiothreitol		7. 0.1% Triton X-100
	1,000 mM NaCl		

Compositions of 10X Loading Buffer (Stored at RT after used):

0.9% SDS
50% Glycerol
0.05% Bromophenol Blue

Add >1/10 volume of 10X Loading Buffer to stop enzyme reaction and apply on agarose gel electrophoresis. SDS may precipitate during the storage at room temperature. In case precipitates generated, dissolve in warm bath before use.

Note

This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals. Also, do not use this product as food, cosmetic, or household item, etc. Takara products may not be resold or transferred, modified for resale or transfer, or used to manufacture commercial products without written approval from Takara Bio Inc. If you require licenses for other use, please contact us by phone at +81 77 565 6973 or from our website at www.takara-bio.com. Your use of this product is also subject to compliance with any applicable licensing requirements described on the product web page. It is your responsibility to review, understand and adhere to any restrictions imposed by such statements. All trademarks are the property of their respective owners. Certain trademarks may not be registered in all jurisdictions.

