


EcoO109I (Dra II)



Code No. 1043A **Size:** **2,000 U**
Conc.: **15 U/ μ l**

Supplied Reagents:

10X L Buffer **1 ml**
10X Loading Buffer **1 ml**

Storage Buffer: 10 mM Tris-HCl, pH 7.5
 400 mM KCl
 0.1 mM EDTA
 1 mM DTT
 0.15% Triton X-100
 0.01% BSA
 50% Glycerol

Storage: -20°C

Source: *Escherichia coli* H709c

General Reaction Mixture:

EcoO109I 1 μ l
 10X L Buffer 2 μ l
 Substrate DNA \leq 1 μ g
 Sterile purified water up to 20 μ l

Reaction Temperature: 37°C

Unit definition:

One unit is defined as the amount of this enzyme required to digest completely 1 μ g of λ DNA in 50 μ l of the reaction mixture at 37°C for 1 hr.

Quality Control Data:

Please see the Certificate of Analysis (CoA) for each lot. You can download the CoA on Takara Bio website.

Relative Activity in Takara Bio's Universal Buffers:

Universal Buffer	L	M	H	K	T (+BSA)
Relative Activity (%)	100	60	<20	<20	100

Ionic Effect on Activity in Basal Buffer:

Salt (mM)	0	20	40	60	80	100	150
NaCl (%)	100	100	120	120	100	60	0
KCl (%)	100	120	160	120	120	80	0

Composition of Basal Buffer:

10 mM Tris-HCl, pH 8.0
 7 mM MgCl₂
 40 mM KCl
 7 mM 2-mercaptoethanol

Number of Cleavage Sites in DNA:

	SV	ϕ X	pBR	pUC	pUC	M13	Col	
λ	Ad2	40	174	322	19	119	mp18	E1
3	44	3	0	4	1	1	0	0

Effect of DNA methylation:

When the sequence including the recognition site is (A/G)GGNCCTGG, the enzyme activity is affected by dcm methylase. In this case, general DNA originated from *E. coli* can not be cleaved by this enzyme.

Compositions of Universal Buffer (Stored at -20°C):

1. 10X L	100 mM Tris-HCl, pH7.5	4. 10X K	200 mM Tris-HCl, pH8.5
	100 mM MgCl ₂		100 mM MgCl ₂
	10 mM Dithiothreitol		10 mM Dithiothreitol
2. 10X M	100 mM Tris-HCl, pH7.5		1,000 mM KCl
	100 mM MgCl ₂	5. 10X T	330 mM Tris-Ac, pH7.9
	10 mM Dithiothreitol	(BSA-free)	100 mM Mg-Ac
	500 mM NaCl		5 mM Dithiothreitol
3. 10X H	500 mM Tris-HCl, pH7.5		660 mM K-Ac
	100 mM MgCl ₂		6. 0.1% BSA
	10 mM Dithiothreitol		7. 0.1% Triton X-100
	1,000 mM NaCl		

Compositions of 10X Loading Buffer (Stored at RT after used):

0.9% SDS
 50% Glycerol
 0.05% Bromophenol Blue

Add >1/10 volume of 10X Loading Buffer to stop enzyme reaction and apply on agarose gel electrophoresis. SDS may precipitate during the storage at room temperature. In case precipitates generated, dissolve in warm bath before use.

Note

This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals. Also, do not use this product as food, cosmetic, or household item, etc. Takara products may not be resold or transferred, modified for resale or transfer, or used to manufacture commercial products without written approval from Takara Bio Inc.

If you require licenses for other use, please contact us by phone at +81 77 565 6973 or from our website at www.takara-bio.com.

Your use of this product is also subject to compliance with any applicable licensing requirements described on the product web page. It is your responsibility to review, understand and adhere to any restrictions imposed by such statements.

All trademarks are the property of their respective owners. Certain trademarks may not be registered in all jurisdictions.

